

## Review Article

## Etiopathogenesis of idiopathic scoliosis

Apostolos M. Papadoudis<sup>1,2</sup>, Kleopatra L. Skourti<sup>3</sup>

<sup>1</sup>Department of Orthopaedics, Athens Naval Hospital, Athens, Greece; <sup>2</sup>Postgraduate Program "Metabolic Bone Diseases", National and Kapodistrian University of Athens, Greece; <sup>3</sup>Department of Pediatrics, 401 Military Hospital of Athens, Athens, Greece

### Abstract

Scoliosis is a three-dimensional deformity of the spine. The key radiographical tool for diagnosis is the Cobb angle and the most common form is a right thoracic convexity with a compensatory left lumbar convexity. Scoliosis is divided, in terms of etiology, into idiopathic, congenital, and secondary. Research into the etiology of idiopathic scoliosis has focused on multiple areas and a great number of suggestions has been made during the last decades concerning hormones, genetic, metabolic and biomechanical factors. Although a lot of theories have been proposed, none is capable to fully describe the pathophysiology of the disease, underlying the complexity and the multifactorial etiology of the condition. The purpose of this review is to summarize the main concepts of etiology and highlight the pathogenetic theories under research in order to reveal possible "targets" for our efforts to understand, diagnose and treat idiopathic scoliosis.

**Keywords:** Idiopathic, Scoliosis, Pathophysiology, Etiology, Pathogenesis

Scoliosis is a three dimensional deformity of the spine characterized by rotation of the vertebrae in the transverse plane and curvatures in the frontal and sagittal plane<sup>1</sup>. The key radiographical tool for diagnosis is Cobb angle, which is the angle created by lines crossing over the upper limit of the first vertebra in the curve and the lower limit of the lowest and final vertebra in the anterior-posterior radiography. Although this cannot entirely describe all spinal abnormalities, it is easy to calculate and to assess and has become the gold standard for scoliotic diagnosis<sup>2</sup>. Deformity is the main clinical sign during the examination, not followed by the presence of pain. The spine may be obviously deviated from the midline, or this may become apparent only when the patient bends forward (the Adams test)<sup>3</sup>. The major curvature may be located at the thoracic, thoracolumbar or lumbar part of the vertebral column and the convexity may point to the left or right side, with compensatory and smaller curvatures above and below. The most common pattern is a thoracic curvature with right convexity with a compensatory left convexity at the lumbar vertebrae<sup>4</sup>.

Scoliosis is divided, in terms of etiology, into idiopathic (no known cause or disease), congenital (present at birth due to congenital etiological factors), and secondary (where it is caused by another disease)<sup>2</sup>. Idiopathic scoliosis consists, moreover, of three sub-categories, according to the age of diagnosis, as infantile (between 0-3 years old), juvenile (between 4-9 years old) and adolescent (age 10

up to maturity)<sup>5</sup>. Adolescent idiopathic scoliosis (AIS) has a prevalence rate between 1-3% and is more common in girls during puberty. The number of girls with AIS is two times higher than the number of boys with this disease and the prevalence is eight times higher among girls than boys when the deformation consists of curvatures greater than 30 degrees<sup>6</sup>.

Research into the etiopathogenesis of idiopathic scoliosis (IS) has spread through the years to multiple areas and a great number of suggestions has been made during the last decades concerning hormones, genetic, metabolic and biomechanical factors. Although a lot of theories have been proposed, none is capable to fully describe the pathophysiology of the disease, underlying the complexity and the multifactorial etiology of the condition. The purpose of this review is to summarize the main concepts of etiology

*The authors have no conflict of interest.*

**Corresponding author:** Apostolos M. Papadoudis, MD, Kottou 23, 14123, Lykovrysi, Athens, Greece

**E-mail:** tolispapad@windowslive.com

**Edited by:** Konstantinos Stathopoulos

**Accepted 15 March 2019**

and highlight the pathogenetic theories under research in order to reveal possible “targets” for our efforts to understand, diagnose and treat idiopathic scoliosis.

## Genetic factors

The inheritance pattern of IS is not fully understood. In 1968, Wynne-Davies studied the familial incidence of the disease in comparison with the general population. They concluded that in 6.9% of first-grade relatives, 3.7% of second-grade relatives and 1.6% of third-grade relatives scoliotic curvatures had been diagnosed. Based on these results, they suggested that genetic factors play a key role in idiopathic scoliosis<sup>7</sup>. A recent study on a Swedish population of monozygotic and dizygotic twins concluded that genetic factors are responsible for the 38% of the scoliotic phenotype, leaving the rest 62% to environmental factors<sup>8</sup>.

Wise et al first described in 2000 DNA regions associated with IS on chromosomes 6, 10 and 18 during a study on an affected seven-member family<sup>9</sup>. Since then, numerous studies have tried to identify loci linked to IS but only the following have all the criteria for statistical significance: 3q12.1, 5q13.3, 6q15-q21, 9q31.2-34.2, 10q23-q25.3, 12p, 17p11, 19p13.3 and Xq22.3-27.2<sup>10</sup>.

Since then, numerous studies have attempted to recognize genes related to connective tissue, bone metabolism and growth, the melatonin pathway and sex hormones. Of the various genes, TIMP2 was linked to curve prognosis in a Chinese population cohort and CALM1, LEP, IL6 and VDR found to be connected with curve predisposition<sup>10</sup>. In a recent study, matrilin 1 gene (MATN1) was examined about its association with AIS. Matrilin 1 is a cartilage matrix protein, has a crucial role in the assembly of the extracellular matrix and is essential for the support of the vertebral column<sup>11</sup>. Interleukin-6 (IL-6) is a proinflammatory cytokine, necessary for the complex cascade that leads to the degeneration of the intervertebral discs. The homozygous GG genotype of IL-6 gene was linked to a higher prevalence of scoliosis in a recent study with Bulgarian patients and the presence of the G allele could be considered as susceptibility factor to IS<sup>12</sup>. The estrogen receptor alpha gene (ESR1) has been tested as a gene connected with IS. ESR1 gene is expressed in both human bone cells (osteoclasts and osteoblasts) and its mutations were recognized as a causative factor for increased bone resorption and impaired skeletal growth. In 2015, a case-control study demonstrated that the ESR1 PvuII genetic polymorphism is associated with IS and curve severity in the Bulgarian population<sup>13</sup>. The vitamin D receptor (VDR) is defined as a ligand-activated transcription factor that controls gene expression in several cell types, including immune cells. Vitamin D-binding protein (VDBP) acts as a plasma protein, serving for vitamin D transport. Wang et al (2016) demonstrated that Bsm I Bb genotype in VDR gene and rs222020 CC genotype in VDBP gene are associated with AIS, and VDR and VDBP genes were correlated with brace treatment efficacy. Moreover, they concluded that

BMI, an essential marker for the prognosis of AIS, is also associated with these polymorphisms<sup>14</sup>. In addition, a recent report suggested that mutation of PTK7 and VANGL1 genes, encoding regulators of WNT/planar cell polarity (PCP) signaling pathway, may be associated with AIS. The authors demonstrated that animal models with a loss of function of these genes developed a spinal malformation resembling to AIS and identified a rare PTK7 missense mutation in a patient with severe AIS<sup>15</sup>. In addition, one of the most important gene polymorphisms discovered so far is the rs11190870 polymorphism of ladybird homeobox 1 (LBX1) gene. LBX1 is crucial for the dorsal spinal neurons and somatosensory system. This variant is suspected for the disease in many populations through abnormal spinal development or neural system growth<sup>16</sup>.

## Role of hormones

Melatonin is the main hormone produced in the pineal gland. Darkness, mainly, stimulates the synthesis and release of melatonin and its presence in human body is essential for circadian rhythm, bone structure, sexual maturation and aging<sup>17</sup>. Part of the physiological function of melatonin is generated through calmodulin and the association between melatonin and calmodulin is featured by high affinity and reversibility<sup>18</sup>. Calmodulin is a receptor protein that binds calcium and, therefore, regulates many cell functions. Melatonin binds, also, to calmodulin and acts as a calmodulin antagonist and, thus, it may modulate diurnally calcium transport<sup>19</sup>. Historically, Dubouset et al were the first to demonstrate that scoliotic curvatures can be diagnosed in pinealectomized chickens if the pinealectomy is performed shortly after hatching. Thus, these data highlight that melatonin deficiency could be responsible for the development of spinal deformities in this animal model, and possibly in humans as well<sup>20</sup>. Genetic studies have showed that variants of the melatonin receptor 1B (MTNR1B) gene are associated with the development of AIS. MTNR1B polymorphism may lead to a down regulation of melatonin receptor 1B expression causing a clinical deficiency of the hormone<sup>21</sup>. Other studies have concluded that there is an abnormal melatonin-signaling pathway in the osteoblasts of AIS patients<sup>22</sup> and that the application of the melatonin antagonist of Luzindole and the calmodulin antagonist of Tamoxifen can improve the success rate of scoliosis in a bipedal rat model<sup>23,24</sup>. However, human melatonin levels are not consistent in different populations and melatonin deficiency as an etiological factor of scoliosis cannot be supported<sup>17</sup>. On the other hand, posture and its related forces on the spine in coordination with melatonin deficiency could play a crucial role for the onset of scoliosis as, theoretically, asymmetry of melatonin release in paravertebral muscles bilaterally could influence the balance of development of paravertebral muscles and could result to scoliosis<sup>1,17</sup>.

Other hormones, which are associated with bone growth and skeletal maturation and IS are vitamin D and sex

hormones. Vitamin D deficiency during first years of life is associated with low mineralization of the bone matrix and can cause a great number of skeletal deformities. Recent studies have demonstrated overall lower levels of vitamin D in patients with AIS, compared with the control groups, indicative of a possible vitamin-D resistance<sup>25</sup>. However, the exact association between Vitamin D and AIS relation remains unclear. It is likely that vitamin D plays a role, either causative or mediating, in the etiopathogenesis of AIS, and therefore larger population-based studies are needed<sup>25,26</sup>. Estrogens are not a causative factor for AIS but they can regulate the severity of spinal curvature. Reduced amounts of LH and FSH released by hypophysis in girls with AIS cause low levels of estrogens, which results in increased osteoclast activity and bone resorption. A defect in the interaction between sex hormones and cellular signal transduction is another factor which can be involved in the etiopathogenesis of scoliosis<sup>6</sup>.

### Central nervous system

A subclinical abnormality of the central nervous system may be an etiological factor for IS. In the 1980s, Herman et al suggested that scoliosis may be the outcome of an alteration in the higher cortical function<sup>27</sup>. In 2011, a study concluded that an impaired pathway in cortical activation during movements in patients with IS is present. The major observation of this study was that patients with IS demonstrate over activation of secondary motor areas at cortical level when compared to healthy controls. An impaired sensorimotor interaction at cortical level can lead to abnormal motor stimuli and imbalance in spine posture causing spine deformity<sup>28</sup>. In addition, other studies have demonstrated motor and somatosensory abnormalities and intra-cortical signaling alterations that are associated with IS<sup>29,30</sup>.

The vestibular system is the sensory system of sound and equilibrium that has also a causative role in the etiopathogenesis of IS. It can produce a neurogenic shift on the vertebral column by activating particular paravertebral muscles bilaterally but asymmetrically. Changes in labyrinth stimuli observed in AIS may contribute to scoliosis through vestibular-spinal interactions. Despite the research proposing a possible etiological role of vestibular system, the evidence linking scoliosis and the vestibular dysfunction are not persistent in all studies<sup>31</sup>.

In 2009, Burwell et al suggested a novel neurogenic theory for AIS pathogenesis, which proposes the imbalance in development between somatic and autonomic nervous systems expressed in the vertebral column producing systemic skeletal overgrowth<sup>32</sup>.

### Biomechanical factors

It has been noted in IS that there is an imbalance between the growth of the vertebral bodies and that of the posterior structures of the column, causing the faster development of the vertebral bodies. The reduced dorsal growth blocks the

increase in height, resulting in rotation of vertebral bodies, in order to create space for themselves. This mechanism produces a rotational lordosis<sup>33</sup>. According to the Hueter-Volkman's law, bone growth in the period of skeletal growth is retarded by compression forces on the growth plate and accelerated by growth plate tension. Because of the curvature in the normal thoracic spine, compressive force is delivered on the ventrally located part of the vertebral column, whereas distractive force is delivered on the dorsally located part<sup>34</sup>. These forces in addition to the rotational lordosis, affecting the spinal segments, could result in a progressive degeneration of individual vertebrae and ultimately lead to progressive curvature of the spine. In addition to the former described mechanism, there is also a three-dimensional deformity of the vertebral bodies itself<sup>33</sup>.

Moreover, other studies have demonstrated that AIS patients have lower lumbar spine bone mineral density (LSBMD), distraction of columns of chondrocytes in the growth plate, deficiency of proteoglycan matrix in the intervertebral disc and low levels of proteins in the paravertebral muscles on the concave side of the curvature compared with the convex paravertebral muscles<sup>35,36</sup>.

### Metabolic factors

In 2014, Clark et al demonstrated metabolic factors present at 10 years old study participants that were linked to scoliosis diagnosed at 15 years of age. These factors include low fat and lean mass and decreased levels of leptin in blood circulation<sup>37</sup>. Leptin is shown to participate in body growth regulation and development especially during puberty<sup>38</sup>. Finally, in 2016, Burwell et al published a paper formulating a new multifactorial concept for the pathogenesis of AIS, unifying the above mentioned theories. The researchers speculated that the decreased circulating leptin is linked to the onset of the imbalanced neuro-osseous growth mechanism that leads to relative anterior skeletal overgrowth. This mechanism in combination with other factors, including anatomical differences of vertebrae, bipedal posture, growth acceleration during puberty, Hueter-Volkman law and decreased bone mass leads to the development of IS<sup>39</sup>.

As far as lifestyle factors are concerned, associations between the onset of scoliosis and participation in sports, such as ballet dancing, rhythmic gymnastics, swimming, and other athletics have also been reported. A recent study demonstrated that AIS is associated with classical ballet training and, especially, frequency of training, number of years of experience, and duration of training in ballet. AIS was also associated with a low body mass index (BMI). However, no association between AIS and passive smoking was observed<sup>40</sup>.

### Conclusions

In spite of the great number of studies conducted, the primary etiology of IS remains unknown and no pathogenetic

theory can fully describe the initiation of spinal curvatures. Instead, idiopathic scoliosis is possible to be a multifactorial disease. Nowadays, treatment is, mainly, focused on the clinical consequences of IS and not on the main reason causing the disease. Despite all these, research on the prognosis of IS would probably be of great interest for our medical practice, in order to minimize X-ray tests, unnecessary surgeries and follow-up visits of patients. Therefore, more research into etiological factors of the disease and large-scale studies in different populations have to be conducted.

## References

- Man GC, Wang WW, Yim AP, Wong JH, Ng TB, Lam TP, Lee SK, Ng BK, Wang CC, Qiu Y, Cheng CY. A Review of pinealectomy-induced melatonin-deficient animal models for the study of etiopathogenesis of adolescent idiopathic scoliosis. *Int J Mol Sci* 2014;15(9):16484-99.
- Goldberg CJ, Moore DP, Fogarty EE, Dowling FE. Scoliosis: a review. *Pediatr Surg Int* 2008;24(2):129-44.
- Solomon L, Warwick D, Nayagam S. *Appley's system of orthopaedics and fractures*. London: Hodder Arnold;2010
- Grauers A, Einarsdottir E, Gerdhem P. Genetics and pathogenesis of idiopathic scoliosis. *Scoliosis Spinal Disord* 2016;11:45.
- Yaman O, Dalbayrak S. Idiopathic scoliosis. *Turk Neurosurg* 2014;24(5):646-57.
- Kulis A, Goździalska A, Drąg J, Jaśkiewicz J, Knapik-Czajka M, Lipik E, Zarzycki D. Participation of sex hormones in multifactorial pathogenesis of adolescent idiopathic scoliosis. *Int Orthop* 2015;39(6):1227-36.
- Wynne-Davies R. Familial (idiopathic) scoliosis. A family survey. *J Bone Joint Surg (Br)* 1968;50(1):24-30.
- Grauers A, Rahman I, Gerdhem P. Heritability of scoliosis. *Eur Spine J* (2012) 21:1069.
- Wise CA, Barnes R, Gillum J, Herring JA, Bowcock AM, Lovett M. Localization of susceptibility to familial idiopathic scoliosis. *Spine* 2000;25(18):2372-80.
- Gorman KF, Julien C, Moreau A. The genetic epidemiology of idiopathic scoliosis. *Eur Spine J* 2012;21(10):1905-19.
- Montanaro L, Parisini P, Greggi T, Di Silvestre M, Campoccia D, Rizzi S, Arciola CR. Evidence of a linkage between matrilin-1 gene (MATN1) and idiopathic scoliosis. *Scoliosis* 2006;1:21.
- Nikolova S, Dikova M, Dikov D, Djerov A, Dzhebir G, Atanasov V, Savov A, Kremensky I. Role of the IL-6 gene in the etiopathogenesis of idiopathic scoliosis. *Anal Cell Pathol (Amst)* 2015;2015:621893.
- Nikolova S, Yablanski V, Vlaev E, Stokov L, Savov A, Kremensky I. Association between estrogen receptor alpha gene polymorphisms and susceptibility to idiopathic scoliosis in Bulgarian patients: A case-control study. *Open Access Maced J Med Sci* 2015;3(2):278-82.
- Wang Y, Cui ZQ, Luo TB, Liu L. Correlations of VDR and VDBP genetic polymorphisms with susceptibility to adolescent idiopathic scoliosis and efficacy of brace treatment. *Genomics* 2016;108(5-6):194-200.
- Andersen MR, Farooq M, Koefoed K, Kjaer KW, Simony A, Christensen ST, Larsen LA. Mutation of the planar cell polarity gene *VANGL1* in adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2016; Publish ahead of Print.
- Zhu Z, Xu L, Qiu Y. Current progress in genetic research of adolescent idiopathic scoliosis. *Ann Transl Med* 2015;3(Suppl 1):S19.
- Girardo M, Bettini N, Dema E, Cervellati S. The role of melatonin in the pathogenesis of adolescent idiopathic scoliosis (AIS). *Eur Spine J* 2011;20(Suppl 1):S68-74.
- Wu JZ, Wu WH, He LJ, Ke QF, Huang L, Dai ZS, Chen Y. Effect of melatonin and calmodulin in an idiopathic scoliosis model. *Biomed Res Int* 2016;2016:8460291.
- Lowe TG, Edgar M, Margulies JY, Miller NH, Raso VJ, Reinker KA, Rivard CH. Etiology of idiopathic scoliosis: current trends in research. *J Bone Joint Surg Am* 2000;82-A(8):1157-68.
- Dubousset J. Experimental scoliosis induced by pineal and diencephalic lesions in young chickens: its relation with clinical findings. *Orthoped Trans* 1983;7:7.
- Yang P, Liu H, Lin J, Yang H. The association of rs4753426 polymorphism in the melatonin receptor 1B (MTNR1B) gene and susceptibility to adolescent idiopathic scoliosis: a systematic review and meta-analysis. *Pain Physician* 2015;18(5):419-31.
- Acaroglu E, Bobe R, Enouf J, Marcucio R, Moldovan F, Moreau A. The metabolic basis of adolescent idiopathic scoliosis: 2011 report of the "metabolic" workgroup of the Fondation Yves Cotrel. *Eur Spine J* 2012;21(6):1033-42.
- Yang S, Zheng C, Jiang J, Lu F, Xia X, Zhu W, Jin X, Ma X. The value of applying a melatonin antagonist (Luzindole) in improving the success rate of the bipedal rat scoliosis model. *BMC Musculoskelet Disord* 2017;18(1):137.
- Akel I, Demirkiran G, Alanay A, Karahan S, Marcucio R, Acaroglu E. The effect of calmodulin antagonists on scoliosis: bipedal C57BL/6 mice model. *Eur Spine J* 2009;18(4):499-505.
- Goździalska A, Jaśkiewicz J, Knapik-Czajka M, Drąg J, Gawlik M, Cieśla M, Kulis A, Zarzycki D, Lipik E. Association of calcium and phosphate balance, vitamin D, PTH, and calcitonin in patients with adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)* 2016;41(8):693-7.
- Balioğlu MB, Aydın C, Kargın D, Albayrak A, Atıcı Y, Tas SK, Kaygusuz MA. Vitamin-D measurement in patients with adolescent idiopathic scoliosis. *J Pediatr Orthop B* 2017;26(1):48-52.
- Herman R, Mixon J, Fisher A, Maulucci R, Stuyck J. Idiopathic scoliosis and the central nervous system: a motor control problem. The Harrington lecture, 1983. Scoliosis Research Society. *Spine (Phila Pa 1976)* 1985;10(1):1-14.
- Domenech J, García-Martí G, Martí-Bonmatí L, Barrios C, Tormos JM, Pascual-Leone A. Abnormal activation of the motor cortical network in idiopathic scoliosis demonstrated by functional MRI. *Eur Spine J* 2011;20(7):1069-78.
- Doménech J, Tormos JM, Barrios C, Pascual-Leone A. Motor cortical hyperexcitability in idiopathic scoliosis: could focal dystonia be a subclinical etiological factor? *Eur Spine J* 2010;19(2):223-30.
- Lo YL, Teo A, Tan YE, Fook-Chong S, Guo CM, Yue WM, Chen J, Tan SB, Lee HW, Dan YF. Motor and somatosensory abnormalities are significant etiological factors for adolescent idiopathic scoliosis. *J Neurol Sci* 2015;359(1-2):117-23.
- Hawasli AH, Hullar TE, Dorward IG. Idiopathic scoliosis and the vestibular system. *Eur Spine J* 2015;24(2):227-33.
- Burwell RG, Aujla RK, Grevitt MP, Dangerfield PH, Moulton A, Randell TL, Anderson SI. Pathogenesis of adolescent idiopathic scoliosis in girls - a double neuro-osseous theory involving disharmony between two nervous systems, somatic and autonomic expressed in the spine and trunk: possible dependency on sympathetic nervous system and hormones with implications for medical therapy. *Scoliosis* 2009;4:24.
- Hefti F. Pathogenesis and biomechanics of adolescent idiopathic scoliosis (AIS). *J Child Orthop* 2013;7(1):17-24.
- Stokes IA. Mechanical effects on skeletal growth. *J Musculoskelet Neuronal Interact* 2002;2(3):277-80.
- Newton Ede MM, Jones SW. Adolescent idiopathic scoliosis: evidence

- for intrinsic factors driving aetiology and progression. *Int Orthop* 2016;40(10):2075-2080.
36. Mannion AF, Meier M, Grob D, Müntener M. Paraspinal muscle fibre type alterations associated with scoliosis: an old problem revisited with new evidence. *Eur Spine J* 1998;7(4):289-93.
37. Clark EM, Taylor HJ, Harding I, Hutchinson J, Nelson I, Deanfield JE, Ness AR, Tobias JH. Association between components of body composition and scoliosis: a prospective cohort study reporting differences identifiable before the onset of scoliosis. *J Bone Miner Res* 2014;29(8):1729-36.
38. Wang WJ, Yeung HY, Chu WC, Tang NL, Lee KM, Qiu Y, Burwell RG, Cheng JC. Top theories for the etiopathogenesis of adolescent idiopathic scoliosis. *J Pediatr Orthop* 2011;31(1 Suppl):S14-27.
39. Burwell RG, Clark EM, Dangerfield PH, Moulton A. Adolescent idiopathic scoliosis (AIS): a multifactorial cascade concept for pathogenesis and embryonic origin. *Scoliosis Spinal Disord* 2016;11:8.
40. Watanabe K, Michikawa T, Yonezawa I, Takaso M, Minami S, Soshi S, Tsuji T, Okada E, Abe K, Takahashi M, Asakura K, Nishiwaki Y, Matsumoto M. Physical activities and lifestyle factors related to adolescent idiopathic scoliosis. *J Bone Joint Surg Am* 2017;99(4):284-294.